## Phenols CHEMets® Kit

**K-8012**/ **R-8012**: 0 - 1 & 0 - 12 ppm

## **Test Procedure**

- 1. Fill the sample cup to the 25 mL mark with the sample to be tested (fig. 1).
- Dissolve the crystals on the tip of the ampoule in the sample by stirring the sample briefly with the ampoule tip (fig. 2).
- Place the CHEMet ampoule, tip first, into the sample cup. Snap the tip. The ampoule will fill leaving a bubble for mixing (fig. 3).
- To mix the ampoule, invert it several times, allowing the bubble to travel from end to end.
- 5. Dry the ampoule. Obtain a test result **1 minute** after snapping the tip.
- 6. Obtain a test result using the appropriate comparator.
  - a.Low Range Comparator (fig. 4):

    Place the ampoule, flat end first, into the comparator. Hold the comparator up toward a source of light and view from the bottom. Rotate the comparator until the best color match is found.




Figure 1



Figure 2



Figure 3



Figure 4

b.High Range Comparator (fig. 5):
Place the ampoule between the color standards until the best color match is found




Figure 5

## **Test Method**

The Phenols CHEMets®1 test kit employs the 4-aminoantipyrine chemistry.  $^{2,3,4}$  In an alkaline solution, phenols react with 4-aminoantipyrine to produce a red colored complex. The color forming reaction is initiated by potassium ferricyanide (tip coating). Test results are expressed in ppm (mg/Liter) "equivalent phenol" as  $C_6H_5OH$ .

Most parasubstituted phenols do not produce a color with this reagent. Ferrous iron causes a blue color which can be eliminated by adding several drops of 1% EDTA to the sample before dissolving the tip coating. Sulfide, in excess of 100 ppm, causes a yellow turbidity. Highly contaminated waste waters may require distillation to separate phenols from nonvolatile impurities.

- $1.\ CHE \textit{Mets is a registered trademark of AquaPhoenix Scientific, LLC U.S.\ Patent\ No.\ 3,634,038}$
- 2. APHA Standard Methods, 14th ed., Method 510 C (1975)
- 3. ASTM D 1783 01, Phenolic Compounds in Water, Test Method B
- 4. EPA Methods for Chemical Analysis of Water and Wastes, method 420.1 (1983)

## **Safety Information**

Read SDS before performing this test procedure. Wear safety glasses and protective gloves.

